



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

Whence $dX = k^2[(y^2 - x^2)dx - 2xydy] \div (x^2 + y^2)^2$ and $dY = k^2[(x^2 - y^2)dy - 2xydx] \div (x^2 + y^2)^2$.

$$\text{Hence } \frac{dX}{dY} = \frac{(x^2 - y^2)dy - 2xydx}{(y^2 - x^2)dx - 2xydy}.$$

Hence, if x' and y' be the co-ordinates of a point on the curve, the tangent at its inverse point may be

$$\text{written } \left(y - \frac{k^2 y'}{x'^2 + y'^2} \right) = \frac{(x'^2 - y'^2)dy' - 2x'y'dx'}{(y'^2 - x'^2)dx' - 2x'y'dy'} \left(x - \frac{k^2 x'}{x'^2 + y'^2} \right).$$

Also solved by G. B. M. ZERR.

PROBLEMS.

30. Proposed by E. W. NICHOLS, Professor of Mathematics in the Virginia Military Institute, Lexington, Virginia.

Given the cardioid $r = a(1 - \cos \theta)$; find the area of its circumscribing square formed by tangents making angles of 45° with its axis.

31. Proposed by E. B. ESCOTT, Ann Arbor, Michigan.

Through a point O on the produced diameter AB of a semicircle draw a secant ORR' , so that the quadrilateral $ABRR'$ inscribed in the semicircle shall be a maximum. Prove that in this case, the projection of RR' on AB is equal in length to the radius of the circle. [*Williamson's Diff. Calculus*, 7th edition, p. 189, Ex. 25.]

MECHANICS.

Conducted by B. F. FINKEL, Kidder, Mo. All contributions to this department should be sent to him.

SOLUTIONS TO PROBLEMS.

12. Proposed by Professor J. F. W. SCHEFFER, A. M., Hagerstown, Maryland.

A horizontal table without weight is supported on three points A , B , and C . A weight W is laid upon the table, at a point G . If $AG = a$, $BG = b$, $CG = c$, $\angle AGB = \beta$, and $\angle AGC = \nu$, find the pressures upon A , B , and C .

I. Solution by F. P. MATZ, M. Sc., Ph. D., Professor of Mathematics and Astronomy in New Windsor College, New Windsor, Maryland, and L. B. FRAKER, Weston, Ohio.

From the problem, we have $AB = \sqrt{(a^2 + b^2 - 2ab \cos \beta)} = m$. Simi-

larly $AC=n$, and $CB=p$. Put $\angle AGD=\phi$, $\angle CGF=\psi$, and $\angle BGE=\omega$; then, obviously, $a \sin \phi + b \sin (\beta - \phi) = m \dots (1)$,

$$c \sin \psi + a \sin (\nu - \psi) = n \dots (2),$$


$$\text{and } b \sin \omega + c \sin [2\pi - (\beta + \nu - \omega)] = p \dots (3).$$

Knowing the values of ϕ , ψ , and ω , from (1), (2), and (3), respectively, we obtain for the pressures upon A , B , and C .

$$P_A = a[\sin \phi + \sin(\nu - \psi)] W \dots (4),$$

$$P_B = b[\sin \omega + \sin(\beta - \phi)] W \dots (5),$$

$$\text{and } P_C = c[\sin \psi - \sin(\beta + \nu - \omega)] W \dots (6).$$



II. Solution by ALFRED HUME, C. E., D. Sc., Professor of Mathematics, University of Mississippi, University P. O., Mississippi, A. H. BELL, Hillsboro, Illinois; and the PROPOSER.

Let the pressures at A , B , and C be represented by these letters.

Taking moments about AG , $b \sin \beta \cdot B = c \sin \gamma \cdot C$.

Taking moments about CG , $a \sin \gamma \cdot A = -b \sin (\beta + \gamma) \cdot B$.

Also, $A + B + C = W$. From these,

$$B = \frac{\sin \gamma}{b} W + \left(\frac{-\sin(\beta + \gamma)}{a} + \frac{\sin \gamma}{b} + \frac{\sin \beta}{c} \right) W.$$

Denoting the quantity in the parenthesis by K , we have, from considerations of symmetry, $A = \frac{-\sin(\beta + \gamma)}{aK} W$, $C = \frac{\sin \beta}{cK} W$.

Excellent solutions to this problem were received from G. B. M. ZERR, and P. H. PHILBRICK

13. Proposed by G. B. M. ZERR, A. M., Principal of High School, Staunton, Virginia.

A man, horse and buggy are going around a circular race course at a 2:40 gait. If the whole outfit weighs 1500 lbs, radius of course is 500 feet and track is inclined so that pressure is equal upon the wheels, find the pressure on the ground due to whole weight.

Solution by the PROPOSER.

In the figure, let C , the bob of the conical pendulum, represent the whole weight of man, horse and buggy $= M$. Let r = radius of curve, V = velocity per second, f = centrifugal force. Then $CG = m$, $CE = f$. Resolve CG into its components CH perpendicular to, and CL parallel to AC ; also, CE into its components $CD = CH$ perpendicular to, and CK parallel to AC .

Then pressure $= P = m \cos \theta + f \sin \theta$, where $\theta = \angle OAC$.

But $f = \frac{mv^2}{gr}$, and $v^2 = \frac{gr^2}{h}$, where $h = AO$, $r = OC$.

$$\sin \theta = \frac{r}{\sqrt{(r^2 + h^2)}}, \cos \theta = \frac{h}{\sqrt{(r^2 + h^2)}}. \therefore P = \frac{mV(r^2 + h^2)}{h}.$$

Now let abc be a section of the road bed; then abc is similar to OAC ,

$ab = a$, $bc = e$, $ac = w$. Then $\frac{\sqrt{(r^2 + h^2)}}{h} = \frac{a}{w} = \frac{a}{a \cos \theta}$. From the triangle